

NEAR-INFRARED SPECTROSCOPY OF SIMPLE ORGANIC MOLECULES IN THE GV TAU N PROTOPLANETARY DISK

ERIKA GIBB, *Physics & Astronomy, University of Missouri St. Louis, St. Louis, MO, USA.*

T Tauri stars are low mass young stars that may serve as analogs to the early solar system. Observations of organic molecules in the protoplanetary disks surrounding T Tauri stars are important for characterizing the chemical and physical processes that lead to planet formation. We used NIRSPEC on Keck 2 to perform a high resolution ($\lambda/\Delta\lambda \sim 25,000$) L-band survey of T Tauri star GV Tau N, a nearly edge-on young star in the L1524 molecular cloud. The nearly edge-on orientation is rare but necessary to sample the disk in absorption, rather than the more common emission line measurements. GV Tau N is one of only two sources for which HCN and C₂H₂ have been reported in absorption (Gibb et al. 2007; Doppmann et al. 2008). More recently, we reported the first detection of methane, CH₄ (Gibb & Horne 2013). The rotational temperatures are relatively high, implying that HCN, C₂H₂, CH₄, and water originate in the warm molecular layer of the inner protoplanetary disk. Differences in rotational temperature for different molecules suggest that the absorbing column for each molecule samples a different radial distribution.

Doppmann, G. W., Najita, K. R., & Carr, J. S. 2008, ApJ, 685, 298

Gibb, E. L., Van Brunt, K. A., Brittain, S. D., & Rettig, T. W. 2007, ApJ, 660, 1572

Gibb, E. L., Horne, D. 2013, ApJ, 776, L28

E.L.G. was supported by NSF Astronomy grant AST-0908230 and NASA Exobiology grant NNX07AK38G.